

Emergency Department Complications During Early Pregnancy

General Information

Vaginal bleeding and abdominal pain in early pregnancy can occur. It is not caused by anything you did—it is not caused by any food, exercise, activity or stress. About one quarter (1 in 4) of pregnant woman who go on to have healthy babies have some bleeding in early pregnancy. Most women who have vaginal bleeding or abdominal discomfort during their pregnancy do not go on to have a miscarriage. However, either can be a warning sign that problems may exist. There is nothing that can be done to stop or prevent a miscarriage that has already begun.

Anytime bleeding or abdominal pain occurs during pregnancy it is important to see a doctor so that appropriate examinations and tests can be carried out. As long as it has been determined that your pregnancy is in the uterus, there are 2 possible outcomes from this complication in early pregnancy. The most likely is that the bleeding or pain will taper off, and bloodwork and ultrasounds done over the next few days and weeks will determine that your pregnancy is progressing normally. The other possibility is that the bleeding or pain may continue or increase, and you may miscarry. Unfortunately, it is usually not possible to determine which of these will happen during your emergency department visit. This can be very frustrating, but it is important to understand that there is nothing that can be done to stop a miscarriage if it is going to occur.

Instructions

Follow up with your family doctor or specialty clinic to ensure the appropriate bloodwork and ultrasounds occur, as per the Emergency doctor's instructions.

When the doctor sends you home:

- Restrict your level of activity according to the doctor's instructions.
- Do not douche or insert any creams or suppositories into the vagina.
- Do not have sexual intercourse until your doctor has indicated that it is safe to do so.
- Do not take baths while you are bleeding - shower instead.
- Do not use tampons while you are bleeding - use sanitary pads instead.

Return to Emergency if any of the following develop:

- The amount of bleeding increases. You need to change your pad more than twice an hour.
- You develop a fever.
- You have pain that is more severe than mild menstrual cramps.
- The pain becomes one sided.
- You feel light-headed or dizzy.

For more information or questions contact the HealthLinkBC at 811.

For more copies, go online at <http://vch.eduhealth.ca> or
email pchem@vch.ca and quote Catalogue No. **GH.200.B62**
© Vancouver Coastal Health, September 2017

The information in this document is intended solely for the
person to whom it was given by the health care team.
www.vch.ca